Requested document: | EP0094564 click here to view the pdf document

Azolyl-methyl oxiranes, their preparation and use as medicaments and plant protecting agents

Patent Number: EP0094564

Publication

date:

1983-11-23

Inventor(s):

JANSSEN BERND DR; MEYER NORBERT DR; POMMER ERNST-HEINRICH DR;

AMMERMANN EBERHARD DR; KOHLMANN FRIEDRICH-WILHELM DR; WESENBERG

WALTER; HEBERLE WOLFGANG DR

Applicant(s):

BASF AG (DE)

Requested

Patent:

FP0094564, A3, B1

Application

Number:

EP19830104412 19830505

Priority Number

(s):

DE19823218129 19820514; DE19823218130 19820514

IPC

C07D405/06; C07D405/14; A01N43/64; A01N43/50; A61K31/41; C07C25/24;

Classification:

C07D303/08

EC

A01N43/50, A01N43/653, C07C17/26B4, C07D303/08, C07D521/00B1E2C,

Classification:

C07D521/00B1E2H

Equivalents:

AU1458883, AU557154, T ES8402833, IL68433, NL9710201, NZ204222

Cited

Documents:

EP0061051; EP0023103

Abstract

1. Claims for the Contracting States: BE CH DE FR GB IT LI NL SE An azolylmethyloxirane of the general formula I see diagramm: EP0094564,P22,F4 where A and B are identical or different and, independently of one another, are naphthyl, biphenyl or phenyl, it being possible for the phenyl radical to be substituted by halogen or nitro, or by alkyl, alkoxy or haloalkyl, each of 1 to 4 carbon atoms, or by phenoxy or phenylsulfonyl, and Z is CH or nitrogen, or a physiologically tolerated or plant-tolerated salt thereof. 1. Claims for the Contracting State: AT A process for the preparation of a azolylmethyloxirane of the general formula I see diagramm: EP0094564,P23,F7 where A and B are identical or different, and independently of one another, are naphthyl, biphenyl or phenyl, it being possible for the phenyl radical to be substituted by halogen or nitro, or by alkyl, alkoxy or haloalkyl, each of 1 to 4 carbon atoms, or by phenoxy or phenylsulfonyl, and Z is CH or nitrogen, or a physiologically tolerated salt thereof, wherein (a) a compound of the formula II see diagramm : EP0094564,P24,F1 where A and B have the above meanings and L is a leaving group which can undergo nucleophilic substitution, is reacted with a compound of the formula III see diagramm: EP0094564,P24,F2 where Me is a hydrogen atom or a metal atom, and Z is CH or nitrogen, or (b) a compound of the formula II, where A and B have the above meanings and L is a hydroxyl group, is reacted with a compound of the formula IV see diagramm : EP0094564,P24,F3 where Z has the above meanings and Y is carbon or sulphur, or (c) a compound of the formula V see diagramm: EP0094564,P24,F4 where Z, A and B have the above meanings, is epoxidized, or (d) a compound of the formula VI see diagramm: EP0094564,P24,F5 where Z and A have the above meanings, is reacted with a compound of the general formula VII see diagramm: EP0094564,P24,F6 where B has the above meanings. R**1 and R**2, which may be identical or different, are each methyl or phenyl, and n is 0 or 1, and, if desired, the compound thus obtained is converted into a salt with a physiologically tolerated acid.

Data supplied from the esp@cenet database - I2